

**Ministry of Health of Ukraine
The National Pirogov Memorial Medical University, Vinnytsia**

The list of medicines for use in responses to the Objective Structured Clinical Exam on Surgery, Pediatric surgery

Rp.: Sol. Ketonolac 3% -1ml

D.t.d. № 10 in amp.

S.: 3 times a day I.M.

Rp.: Sol. Diclofenaci 2,5% - 3 ml

D.t.d. №10 in amp.

S.: Once a day I.M.

Rp.: Sol. Nalbuphini 1% 1 ml

D.t.d № 5 in ampullis

S.: 2 times a day I.M.

Rp.: Sol. Morphini hydrochloridi 1% -1 ml

D.t.d. № 10 in amp.

S.: For I.V., I.M. administration every 6 hrs.

Rp: Sol..Metronidazoli 0,5% -100 ml

D.S.: 3 times a day I.V.

Rp: Meropenem 1.0 №10

D.S.: 3 times a day I.V. after dissolving in 100 ml of saline sol.

Rp: Ceftriaxoni 1,0

D. t. d. №10

S.: 2 times a day I.V. after dissolving in 100 ml of saline sol.

Rp.: Cefepimi 1,0

D.t.d №10

S. 2 times a day I.V. after dissolving in 100 ml of saline sol.

Rp.: Sol.Amikacini 25% - 2 ml

D.t.d № 10 in ampullis

S. 2 times a day I.V. after dissolving in 100 ml of saline sol.

Rp: Ciprofloxacini 0,2 №10

D. S.: 2 times a day I.V.

Rp: Laevofloxacini 5% 100 ml N10

D.S.: Once a day I.V.

Rp: Amoxycillini/Klavulanati 0.875/0.125 №10

D.S.: 3 times a day I.V. after dissolving in 100 ml of saline sol.

Rp: Sol. Hydrocortizoni 2,5% 2ml

D.t.d.№10 in amp.

S.: For I.V., I.M. administration

Rp.: Sol. Prednisoloni 3% 1ml

D.t.d. № 10 in amp.
S.: For I.V., I.M. administration

Rp.: Sol. Heparini 5 ml (1 ml-5000 MO)
D.t.d. № 1
S.: For I.V., administration every 6 hrs. under control of blood clotting time

Rp: Sol. Enoxaparini 10% 0,4 ml
D.t.d. № 6 in amp.
S.: Once a day 0,4 ml subcutaneously

Rp.: Sol. Lidocaini hydrochloridi 2% - 2ml
D.t.d. №10 in amp.
S.: For local anesthesia.

Rp.: Pantoprazoli 0,02 №5
D.S.: 2 times a day I.V. after dissolving in 100 ml of saline sol.

Rp.: Sol. Glucosae 5% - 200 ml
D.S.: For I.V. administration

Rp.: Sol. Natrii chloridi 0,9% - 200 ml
D.S.: For I.V. administration

Rp.: Sol. Adrenalini hydrochloridi 0,1% -1ml
D.t.d. № 10 in amp.
S.: For I.V. administration in saline sol.

Rp.: Sol. Noradrenalini hydrotartratis 0.2%-1 ml
D.t.d. № 10 in amp.
S: For I.V. administration in 500 ml of saline sol.

Rp.: Sol. Dobutamini 0.5% -50ml
D.t.d. № 10 in amp.
S.: For I.V. administration in 500 ml of saline sol. 2,5-10 mkg/kg

Rp.: Sol. Amiodaroni 5% 3ml
D.t.d. №.10 in amp.
S.: For I.V. administration in 500 ml of saline sol.

Rp.: Sol. Natrii oxybutiratis 20% -10 ml
D.t.d. № 10 in amp.
S.: For I.V. administration 50 -120 mkg/kg

Rp.: Sol. Atropini sulfatis 0.1% -1 ml
D.t.d. № 10 in amp.
S.: For I.V. administration in saline sol.

Rp.: Sol. Furosemidi 1% -2ml
D.t.d. № 10 in amp.
S.: For I.V. administration in saline sol.

Rp.: Ac. tranexamici 0.5 (5 ml) in amp. №5
S.: 2 times a day I.V.

Rp.: Fresh frozen plasma 200 ml

D.S.: For I.V. transfusion after thawing

Rp.: Red cell mass 200 ml

D.S.: For I.V. transfusion after tests on a group and Rh identity, individual and biological compatibility